arXiv:0811.3398v1 [math.AG] 20 Nov 2008

RATIONALLY CONNECTED FOLIATIONS ON SURFACES

SEBASTIAN NEUMANN

Abstract. In this short note we study foliations with rationally connected leaves on surfaces. Our main result is that on surfaces there exists a polarisation such that the Harder-Narasimhan filtration of the tangent bundle with respect to this polarisation yields the maximal rationally quotient of the surface.

1. Introduction

Let X be a smooth projective variety. In this note, we are interested in foliations with rationally connected leaves. In [KSCT07] the authors show how to construct such foliations from the *Harder-Narasimhan Filtration* of the tangent bundle of the variety. This construction heavily depends on a chosen polarisation, and therefore the question arises how this foliation varies changing the polarisation.

There is another way to construct a fibration with rationally connected fibers, the *maximal rationally connected quotient*. This is a rational map, such that the fibers are rationally connected. Almost every rational curve lies in a fiber of this map.

We can ask if the Harder-Narasimhan filtration of the tangent bundle always induces the maximal rationally quotient with respect to any polarisation. The answer is negative already on surfaces due to an example of T. Eckl ([Eck08]).

In this note we will prove that on surfaces there exists always a polarisation, such that the Harder-Narasimhan filtration yields the maximal rationally quotient.

1.1. **Acknowledgements.** The author was supported in part by the Graduiertenkolleg “Globale Strukturen in Geometrie und Analysis” of the Deutsche Forschungsgemeinschaft. The result is part of the author’s forthcoming Ph.D. thesis written under the supervision of Stefan Kebekus. He would like to thank Stefan Kebekus, Thomas Eckl and Sammy Barkowski for numerous discussions.

The results in this note were presented at a Workshop in Grenoble in April 2008. Seemingly similar results have been obtained independently in [SCT08].

Date: 18.11.2008.

2. PRELIMINARY RESULTS AND NOTATION

Let X be an n-dimensional projective variety over the complex numbers with an ample line bundle H. Given a torsion-free coherent sheaf $\mathcal{F}$ on X, we define the *slope of* $\mathcal{F}$ *with respect to* H by

$$\mu_H(\mathcal{F}) := \frac{c_1(\mathcal{F}).H^{n-1}}{\mathrm{rk}(\mathcal{F})}.$$

We call $\mathcal{F}$ *semistable with respect to* H if for any nonzero proper subsheaf $\mathcal{G}$ of $\mathcal{F}$ we have $\mu_H(\mathcal{G}) \leq \mu_H(\mathcal{F})$.
If there exists a nonzero subsheaf $\mathcal{G} \subset \mathcal{F}$ such that $\mu_H(\mathcal{G}) > \mu_H(\mathcal{F})$, we will call $\mathcal{G}$ a *destabilizing subsheaf* of $\mathcal{F}$.

For a proof of the following result, we refer the reader to [HL97, Theorem 1.3.4].

Theorem 2.1. *Let* $\mathcal{F}$ *be a torsion-free coherent sheaf on a smooth projective variety and* H *be an ample line bundle on* X. *There exists a unique filtration of* $\mathcal{F}$*, the so called* Harder-Narasimhan filtration *or* HN-filtration*, depending on the chosen ample line bundle*

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \ldots \subset \mathcal{F}_k = \mathcal{F}$$

with the following properties:

(i) *The quotients* $\mathcal{G}_i := \mathcal{F}_i/\mathcal{F}_{i-1}$ *are torsion-free and semistable.*
(ii) *The slopes of quotients satisfy* $\mu_H(\mathcal{G}_1) > \ldots > \mu_H(\mathcal{G}_k)$.
(iii) *The sheaves* $\mathcal{F}_i$ *are saturated in* $\mathcal{F}$.

Definition 2.2. Let $\mathcal{F}$ be a torsion-free coherent sheaf on a smooth projective variety. The unique sheaf $\mathcal{F}_1$ appearing in the Harder-Narasimhan filtration of $\mathcal{F}$ is called *the maximal destabilizing subsheaf of* $\mathcal{F}$.

Definition 2.3. Let $\mathcal{F}$ be a coherent torsion-free sheaf on a smooth projective variety with HNF with respect to an ample line bundle H

$$0 = \mathcal{F}_0 \subset \ldots \subset \mathcal{F}_k = \mathcal{F}.$$

If the slope of the quotient $\mathcal{F}_i/\mathcal{F}_{i-1}$ is positive with respect to H, then $\mathcal{F}_i$ is called *positive with respect to* H.

Remark 2.4. Note that the construction of the HNF naturally extends to $\mathbb{Q}$– and $\mathbb{R}$–divisors, i.e we do not need to assume the chosen polarisation to be integral.

Obviously, the Harder-Narasimhan filtration depends only on the numerical class of the chosen ample bundle. In particular it makes sense to ask how the filtration of a given sheaf depends on the ample bundle sitting in the finite dimensional vectorspace of all divisors modulo numerical equivalence.

Notation 2.5. We sometimes omit the polarisation in the notation of the slope, that is we write $\mu(\mathcal{F})$ for the slope of a sheaf $\mathcal{F}$ with respect to a polarisation.

We can now state an important result originally formulated by Miyaoka and explicitely shown in [KSCT07]. For a survey on these and related results we refer the reader to [KSC06].

Theorem 2.6. [KSCT07, Theorem 1] *Let X be a smooth projective variety and let*

$$0 = \mathcal{F}_0 \subset \mathcal{F}_1 \subset \ldots \subset \mathcal{F}_k = TX$$

be the Harder-Narsimhan filtration of the tangent bundle with respect to a polarisation H. Write $\mu_i := \mu_H(\mathcal{F}_i/\mathcal{F}_{i-1})$ for the slopes of the quotients. Assume $\mu_1 > 0$ and set $m := \max\{i \in \mathbb{N} | \mu_i > 0\}$. Then each $\mathcal{F}_i$ with $i \leq m$ is a foliation with algebraic leaves and for general $x \in X$ the closure of the leaf through x is rationally connected.

Let X be a smooth projective variety and assume the conditions of Theorem (2.6) are fulfilled. Thus we obtain foliations $\mathcal{F}_1, \ldots, \mathcal{F}_k$ with algebraic and rationally connected leaves. By setting

$$\begin{array}{rccl} q_i: & X & \dashrightarrow & \mathrm{Im}(q_i) \subset \mathrm{Chow}(X) \\ & x & \mapsto & \mathcal{F}_i\text{-leaf through } x \end{array}$$

we obtain a map, such that the closure of the general fibre is rationally connected, see [KSCT07], section 7.
There is another map with this property called the *maximal rationally connected quotient* or *MRC-quotient* for short based on a construction by Campana [Cam81], [Cam94] and Kollár-Miyaoka-Mori [KMM92].

Theorem 2.7. [Kol96, Chapter IV, Theorem 5.2] *Let X be a smooth projective variety. There exists a variety Z and a rational map $\phi : X \dashrightarrow Z$ with the following properties:*

- *the fibers of ϕ are rationally connected,*
- *a very general fiber of ϕ is an equivalence class under rationally connectivity and*
- *up to birational equivalence the map ϕ and the variety Z are unique.*

It is possible to ask if the HNF with respect to a certain polarisation yields the MRC-quotient. We will give a positive answer in the next section.

3. Rationally connected fibration on surfaces and the MRC-quotient

In this section X will be a smooth projective surface over the complex numbers. We want to investigate the regions in the ample cone which induce the same HN-filtration. More precisely we divide the ample cone into parts, such that in each part, we get the same HN-filtration of the tangent bundle. With this at hand we are able to show that the MRC-quotient comes from the Harder-Narasimhan filtration with respect to a certain polarisation.

In order to compute the HN-filtration of the tangent bundle on surfaces, we only have to search for a destabilizing subbundle, such that the quotient is torsion-free.

Lemma 3.1. *Let X be a smooth projective surface. If $\mathcal{F} \subset TX$ is a destabilizing subsheaf with respect to a polarisation such that $TX/\mathcal{F}$ is torsion-free, then the Harder-Narasimhan filtration is given by $0 \subset \mathcal{F} \subset TX$.*

Proof. Consider the exact sequence

$$0 \to \mathcal{F} \to TX \to TX/\mathcal{F} \to 0.$$

Using that the rank and the chern class is additiv in short exact sequences, we obtain

$$\mu(TX) = \frac{1}{2}\mu(TX/\mathcal{F}) + \frac{1}{2}\mu(\mathcal{F}).$$

Since $\mu(\mathcal{F}) > \mu(TX)$, we therefore have $\mu(\mathcal{F}) > \mu(TX/\mathcal{F})$. That is,

$$0 \subset \mathcal{F} \subset TX$$

satisfies the properties of the Harder-Narasimhan filtration and by the uniqueness of the HN-filtration, we finish the proof. ☐

Notation 3.2. We will write $N^1(X)$ for the Néron-Severi group and $\mathrm{N}^1_{\mathbb{Q}}(X)$ (resp. $N^1_{\mathbb{R}}(X)$) for the vectorspace of $\mathbb{Q}$–divisors (resp. $\mathbb{R}$–divisors) modulo numerical equivalence on X. The convex cone of all ample $\mathbb{R}$–divisors in $N^1_{\mathbb{R}}(X)$ will be denoted by $\mathrm{Amp}_{\mathbb{R}}(X)$.

Now we denote the regions in $\mathrm{Amp}_{\mathbb{R}}(X)$, we are interested in. Let $H \in N^1_{\mathbb{R}}(X)$ be an ample bundle. If TX is not semistable with respect to H, let $D_{TX}(H)$ be the unique maximally destabilizing subbundle of TX. In this case, we call

$$\Delta_H := \left\{\tilde{H} \in \mathrm{Amp}_{\mathbb{R}}(X) | D_{TX}(H) = D_{TX}(\tilde{H})\right\}$$

the *destabilisation chamber with respect to H*. Note that if the tangent bundle is semistable with respect to a certain polarisation, then we get a chamber, such that for all polarisations in this chamber TX is semistable. This region will be denoted by Σ and we call it the *semistable chamber*. That is

$$\Sigma := \{H \in \mathrm{Amp}_{\mathbb{R}}(X) | TX \text{ is semistable with respect to } H\}.$$

The destabilizing chambers and the chamber of stability give a decomposition of the ample cone. Concerning the structure of these chambers we prove the following lemma.

Lemma 3.3. *Let X be a smooth projective surface. We have*

(i) *The destabilisation chambers and the semistable chamber are convex cones in $\mathrm{Amp}_{\mathbb{R}}(X)$.*

(ii) *The semistable chamber is closed in* $\mathrm{Amp}_{\mathbb{R}}(X)$.
(iii) *The destabilisation chambers are open in* $\mathrm{Amp}_{\mathbb{R}}(X)$.

Proof. The convexity property follows from the linearity of the intersection product and the uniqueness of the HN-filtration. In detail: Let H_1 and H_2 be two polarisations belonging to the same chamber. This means, that there exists a subsheaf $L \subset TX$ which is maximal destabilizing both with respect to H_1 and H_2. We obtain

$$c_1(L).(H_1 + H_2) = c_1(L).H_1 + c_1(L).H_2 > \frac{1}{2}c_1(TX).(H_1 + H_2).$$

By Lemma (3.1), we deduce that L is the maximally destabilizing subsheaf with respect to $H_1 + H_2$.
In the same spirit we prove the convexity of Σ: Let H_1 and $H_2 \in \Sigma$. This means by definition that for all subsheaves $L \subset TX$ the inequality $c_1(L).H_i \leq \frac{1}{2}c_1(TX).H_i$, $i = 1, 2$ holds. Therefore by linearity of the intersection product the inequality holds for $H_1 + H_2$. This proves (i).
The second statement follows from the continuity of the intersection product. Let $H_n \in \Sigma$ be a sequence converging to $H \in \mathrm{Amp}(X)$. Then for a given subsheaf $L \subset TX$ one has $\mu_{H_n}(L) \leq \mu_{H_n}(TX)$ for each $n \in \mathbb{N}$. Passing to the limit the inequality still holds by continuity.
Statement (iii) is also a direct consequence of the continuity of the intersection product: Let $H \in \Delta_{\tilde{H}}$. Let L be the maximal destabilizing subsheaf of the tangent bundle. That is we have

$$(c_1(L) - \frac{1}{2}c_1(TX)).H > 0.$$

This inequality holds in a neighbourhood of H. By Lemma (3.1) the subsheaf L gives the HN-filtration in a neighbourhood of H. □

In order to prove our main result, we use the following corollary.

Corollary 3.4. *Let X be a smooth projective surface. Let l be a linesegment in* $\mathrm{Amp}_{\mathbb{R}}(X)$. *If TX is not semistable with respect to all polarisation on l, then l lies completely in one destabilizing chamber.*

In order to prove semistability of the tangent bundle on certain surfaces having many automorphisms, we give a useful lemma.

Lemma 3.5. *Let X be a smooth projective surface and let $\sigma \in \mathrm{Aut}^0(X)$. Let $\mathcal{F}$ be the maximal destabilizing subsheaf of TX with respect to a polarisation. Then we have $\sigma^*(c_1(\mathcal{F})) = c_1(\mathcal{F})$. In particular: If the slope of $\mathcal{F}$ is positive then the automorphism σ maps each leaf of $\mathcal{F}$ to another leaf of $\mathcal{F}$.*

Proof. Let $H \in N^1(X)$ and let $\mathcal{F}$ be the maximal destabilizing subsheaf of TX with respect to H. We compute the slope of $\sigma^*(T_{\mathcal{F}})$:

$$\begin{array}{rcl} \mu(\sigma^*(c_1(\mathcal{F}))) & = & H.\sigma^*(c_1(\mathcal{F})) \\ & = & \sigma_*(H).c_1(\mathcal{F}) \\ & = & H.c_1(\mathcal{F}) \\ & > & \frac{1}{2}c_1(TX).H. \end{array}$$

We give an explanation of the third equality: Recall that the group of automorphisms acts on the Néron-Severi group. Since $N^1(X)$ is discrete $\mathrm{Aut}^0(X)$ acts trivially on $N^1(X)$, i.e. the pushforward of σ has to give the same element in $N^1(X)$.
Therefore, we have shown that $\sigma^*(T_{\mathcal{F}})$ is also a destabilizing bundle and by the uniqueness of the HN-filtration we finish the proof. ☐

Example 3.6. *Hirzebruch Surfaces*
Let Σ_n be the n-th Hirzebruch surface, $\pi : \Sigma_n \longrightarrow \mathbb{P}^1$ the projection onto the projective line. We denote the fiber under the projection with f and the distinguished section with selfintersection $-n$ with C_0. Recall (see [Har77], chapter V.2) that $\mathrm{Num}_{\mathbb{R}}(\Sigma_n) =< C_0, f >$ and a divisor $D \equiv_{num} aC_0 + bf$ is ample if and only if $a > 0$ and $b > an$. The canonical bundle is given by $-K_{\Sigma_n} = 2C_0 + (2 + n)f$. The relative tangent bundle of π is a natural candidate for a destabilizing subbundle. We have the sequence

$$0 \to T_{\Sigma_n/\mathbb{P}^1} \to T\Sigma_n \to \pi^*T\mathbb{P}^1 \to 0$$

Let $H := xC_0 + yf$ be a polarisation. Then one can compute that $T_{\Sigma_n/\mathbb{P}^1}$ is destabilizing if and only if $-2x - nx + 2y > 0$. In particular we compute for $n \geq 2$:

$$-2x - nx + 2y > -2x - nx + 2nx = -2x + nx \geq 0.$$

Therefore, for $n \geq 2$ the HN-filtration is given by

$$0 \subset T_{X/\mathbb{P}^1} \subset TX$$

for all polarisations. In other words we obtain only one destabilizing chamber.

For $n = 0$ we have $\Sigma_0 = \mathbb{P}^1 \times \mathbb{P}^1$ and we get three chambers. Two destabilizing chambers correspond to the two relative tangent bundles of the projections. They are cut out by the inequalitys $x > y$ and $x < y$. There is a chamber of semistability, which is determined by the equation $x = y$.

For $n = 1$ we see that for $x > \frac{3}{2}y$ the relative tangent bundle is destabilizing. Since Σ_1 is the projective plane with one point blown up, say p, we know the group of automorphisms, which is the automorphism group of the projective plane leaving p fixed. The destabilizing foliation corresponds to the radial foliation through p in the plane.
So if there were another foliation $\mathcal{F}$ coming from the Harder-Narasimhan filtration, we could deform the leaves with these automorphisms. Then we


y
$0 \subset T^1_{X/\mathbb{P}^1} \subset TX$
$0 \subset T^2_{X/\mathbb{P}^1} \subset TX$
x

FIGURE 1. The ample cone of $X = \Sigma_0$ and the chamber structure. Here $T^1_{X/\mathbb{P}^1}$ and $T^2_{X/\mathbb{P}^1}$ denote the relative tangent bundle of the first and second projection.

would again obtain leaves of this foliation by Lemma (3.5). So unless $\mathcal{F}$ is the radial foliation, we could produce infintely many singularities, which is absurd. Therefore we conclude that the tangent bundle is semistable for $x \leq \frac{3}{2}y$.

Remark 3.7. From this example follows, that the stabilizing chamber can be both lower dimensional or equal the dimension of the ample cone.

Now we want to answer the question if there always exists a polarisation, such that the HNF gives rise to the MRC-quotient.

Theorem 3.8. *Let X be a uniruled projective surface. Then there exists a polarisation, such that the maximal rationally quotient of X is given by the foliation associated to highest positive term in the HNF associated with this polarisation.*

Proof. To start, observe that there is always a polarisation H_1 such that $c_1(TX).H_1 > 0$. Assume on the contrary, that each ample bundle intersects $c_1(TX)$ negatively or zero. This means that K_X is pseudoeffective. On the other hand there exists a free rational curve $f : \mathbb{P}^1 \to X$. See [Deb01, Corollary 4.11] for a prove of the existence of such a curve. Writing

$$f^*(TX) = \mathcal{O}(a_1) \oplus \mathcal{O}(a_2)$$

with $a_1 + a_2 \geq 2$, we compute

$$K_X.f_*\mathbb{P}^1 = -a_1 - a_2 \leq -2,$$

a contradiction.

First let us assume that X is not rationally connected. Then we take the polarisation H_1 with $c_1(TX).H_1 > 0$. There exists a destabilizing subsheaf $\mathcal{F}$ of TX, since otherwise X would be rationally connected by Theorem (2.6). Furthermore the slope of $\mathcal{F}$ has to be bigger than $c_1(TX).H_1$ and therefore positive. So this sheaf will give a foliation with rationally connected leaves and hence the maximal rationally quotient.

Consider the case where X is rationally connected. Then we fix a very free rational curve l on X. For a proof of the existence of a very free rational curve see [Deb01, Corollary 4.17]. This means that $TX|_l$ is ample. So we know that each quotient of $TX|_l$ has strictly positive degree.
Since l is movable, it is in particular nef. Let H_2 be an ample class. Because l is nef, we know that $H_\epsilon := l + \epsilon H_2$ is ample in $N^1_{\mathbb{Q}}(X)$ for any $\epsilon > 0$. Observe that $c_1(TX).H_\epsilon > 0$ for sufficiently small ϵ, say for $0 < \epsilon < 1$. If TX is semistable with respect to a certain polarisation H_ϵ with $0 < \epsilon < 1$, the claim follows since TX has positive slope and induces a trivial foliation which gives the rationally connected quotient. If TX is not semistable for all polarisations H_ϵ with $0 < \epsilon < 1$, let L_ϵ be the destabilizing subsheaf of TX with respect to H_ϵ. Because of Corollary (3.4) the ray H_ϵ stays in one destablizing chamber. So $L := L_\epsilon$ remains constant.
Now it is clear that for sufficiently small ϵ both, the slope of L and the slope of TX/L will be positive with respect to H_ϵ. Therefore the HN-fitration of TX with respect to H_ϵ yields the maximal rationally connected quotient. $\square$

REFERENCES

[Cam81] F. Campana. Coréduction algébrique d'un espace analytique faiblement kählérien compact. *Invent. Math.*, 63(2):187–223, 1981.

[Cam94] Frédéric Campana. Remarques sur le revêtement universel des variétés kählériennes compactes. *Bull. Soc. Math. France*, 122(2):255–284, 1994.

[Deb01] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.

[Eck08] Thomas Eckl. Lower bounds for Seshadri constants. *Math. Nachr.*, 281(8):1119–1128, 2008.

[Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.

[HL97] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997.

[KMM92] János Kollár, Yoichi Miyaoka, and Shigefumi Mori. Rationally connected varieties. *J. Algebraic Geom.*, 1(3):429–448, 1992.

[Kol96] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996.

[KSC06] Stefan Kebekus and Luis Solá Conde. Existence of rational curves on algebraic varieties, minimal rational tangents, and applications. In *Global aspects of complex geometry*, pages 359–416. Springer, Berlin, 2006.

[KSCT07] Stefan Kebekus, Luis Solá Conde, and Matei Toma. Rationally connected foliations after Bogomolov and McQuillan. *J. Algebraic Geom.*, 16(1):65–81, 2007.

[SCT08] Luis Solá Conde and Matei Toma. Maximally rationally connectes fibrations and movable curves. *preprint, arXiv:0811.2141v2 [math.AG]*, 2008.

E-mail address: `sebastian.neumann@math.uni-freiburg.de`